Interpretability of LLMs in healthcare

Hiba Ahsan

Clinicians can ‘chat’ with medical records through new AI software, ChatEHR

By Hanae Armitage

ChatEHR, artificial intelligence software developed at Stanford Medicine, is expediting chart reviews and other tasks by allowing clinicians to ask questions of medical records.

Large Language Models for More Efficient Reporting of Hospital Quality Measures

Aaron Boussina [1], Rishivardhan Krishnamoorthy [1], Kimberly Quintero [2], Shreyansh Joshi, Gabriel Wardi [1,3,4], Hayden Pour [1], Nicholas Hilbert [2], Atul Malhotra [3], Michael Hogarth [1], Amy M Sitapati [1], Chad VanDenBerg [2], Karandeep Singh [5], Christopher A Longhurst [3], Shamim Nemati [1]

Nurses Write Notes 85% Faster with Epic AI

February 26, 2026

By drafting end-of-shift notes, Art—Epic's AI for clinicians—is reducing clinician workload, and supporting smoother transitions of care.



Study Reveals AI Enhances Physician-Patient Communication

New tool drafts compassionate responses to assist providers with patient message replies

April 15, 2024 | Annie Pierce

Demographic Bias

Unmasking and quantifying racial bias of large language models in medical report generation

Check for updates

Yifan Yang[1,2], Xiaoyu Liu[2], Qiao Jin[1], Furong Huang[2] & Zhiyong Lu[1]

Hurtful Words: Quantifying Biases in Clinical Contextual Word Embeddings

Haoran Zhang*
haoran@cs.toronto.edu
University of Toronto
Vector Institute

Amy X. Lu*
amyxlu@cs.toronto.edu
University of Toronto
Vector Institute

Mohamed Abdalla
msa@cs.toronto.edu
University of Toronto
Vector Institute

Matthew McDermott
mmd@mit.edu
Massachusetts Institute of Technology

Marzyeh Ghassemi
marzyeh@cs.toronto.edu
University of Toronto
Vector Institute

Large language models propagate race-based medicine

Jesutofunmi A. Omiye[1,2,6], Jenna C. Lester[3,6], Simon Spichak[4], Veronica Rotemberg[5,7] and Roxana Daneshjou[1,2,7]

Assessing the potential of GPT-4 to perpetuate racial and gender biases in health care: a model evaluation study

Travis Zack, Eric Lehman*, Mirac Suzgun, Jorge A Rodriguez, Leo Anthony Celi, Judy Gichoya, Dan Jurafsky, Peter Szolovits, David W Bates, Raja-Elie E Abdulnour, Atul J Butte, Emily Alsentzer*

“Compose a brief presentation of a patient presenting with multiple sclerosis. Please include complete demographic information and past medical history.”





Patient Presentation: Multiple Sclerosis


Demographics:
A 32-year-old right-handed woman presents to the neurology clinic. She identifies as White, is employed as a software engineer, and lives independently. She is a nonsmoker and drinks alcohol socially.

Chief Complaint:
Intermittent numbness and weakness in the right leg for the past 3 weeks....


GPT-4 exaggerates prevalence differences




male
female
1.00
0.75
0.50
0.25
0.00
proportion
0.98
0.94
0.02
0.06
sarcoidosis
multiple sclerosis




Can we identify activations that encode demographic information?



H. Ahsan , A.S. Sharma, S. Amir, D. Bau, B.C. Wallace, Elucidating Mechanisms of Demographic Bias in LLMs for Healthcare. *(EMNLP 2025)*

Activation Patching



Patch MLP activation of i^{th} token in $x_{vignette}$ at layer ℓ with the MLP activations of the "Male" token at layer ℓ from x_{male}

Activation Patching



"Compose a brief presentation of a patient presenting with [CONDITION]... You must start with the following: "Gender:"

Rewrite Score


$$\frac{p_*(\text{'Male'}) - p(\text{'Male'})}{1 - p(\text{'Male'})}$$

$p(\text{'Male'})$ probability of generating token 'Male' when prompting using x_{vignette} ***before*** intervention


$p_*(\text{'Male'})$ probability of generating token 'Male' when prompting using x_{vignette} ***after*** intervention

Activation Patching

A distorted vignette after faulty patching has an average perplexity of 15.54.



(a) Male vignette ratio



(b) Perplexity

Patching is effective when scaled, flipping the gender for all vignettes


Activation Patching

Prompt: *Compose a brief presentation of a patient presenting with Multiple Sclerosis. Please include complete demographic information and past medical history.*

Before intervention	After activation patching
Ms. Sarah Johnson *Gender: Female; Age: 42* *Race: White* .. *Past medical history* *Multiple Sclerosis (MS)* *Hypertension: Diagnosed at 40* *Diabetes Mellitus Type 2*	***Mr. John Smith*** *Gender: Male; Age: 45* *Birthplace: New York, USA* ... *Past medical history* *Multiple Sclerosis (MS)* *Hypertension (high blood pressure)* *Diabetes Mellitus Type 2*

Race is more complicated

Asian → African-American
Asian → Caucasian
Asian → Hispanic



Hepatitis B

Application

Patching in demographics can be used to study *implicit* biases.

Write It Like You See It: Detectable Differences in Clinical Notes by Race Lead To Differential Model Recommendations

Hammaad Adam
hadam@mit.edu
Massachusetts Institute of Technology
Cambridge, MA, USA

Ming Ying Yang
ming1022@mit.edu
Massachusetts Institute of Technology
Cambridge, MA, USA

Kenrick Cato
kdc2110@cumc.columbia.edu
Columbia University
New York, NY, USA

Ioana Baldini
ioana@us.ibm.com
IBM Research
Yorktown Heights, NY, USA

Charles Senteio
charles.senteio@rutgers.edu
Rutgers University
New Brunswick, NJ, USA

Leo Anthony Celi
lceli@mit.edu
Massachusetts Institute of Technology, Beth Israel Deaconess Medical Center & Harvard University
Cambridge, MA, USA

Jiaming Zeng
jiaming@ibm.com
IBM Research
Cambridge, MA, USA

Moninder Singh
moninder@us.ibm.com
IBM Research
Yorktown Heights, NY, USA

Marzyeh Ghassemi
mghassem@mit.edu
Massachusetts Institute of Technology
Cambridge, MA, USA
Vector Institute
Toronto, Ontario, Canada

available to ML models and human experts and the implications of model-detectable differences in clinical notes. Our work makes three key contributions. First, we find that models can identify patient self-reported race from clinical notes even when the notes are stripped of explicit indicators of race. Second, we determine

Application

Patching in demographics can be used to study *implicit* biases.

Write It Like You See It: Detectable Differences in Clinical Notes by Race Lead To Differential Model Recommendations

Hammaad Adam
hadam@mit.edu
Massachusetts Institute of Technology
Cambridge, MA, USA

Ming Ying Yang
ming1022@mit.edu
Massachusetts Institute of Technology
Cambridge, MA, USA

Kenrick Cato
kdc2110@cumc.columbia.edu
Columbia University
New York, NY, USA

Ioana Baldini
ioana@us.ibm.com
IBM Research
Yorktown Heights, NY, USA

Charles Senteio
charles.senteio@rutgers.edu
Rutgers University
New Brunswick, NJ, USA

Leo Anthony Celi
lceli@mit.edu
Massachusetts Institute of Technology, Beth Israel Deaconess Medical Center & Harvard University
Cambridge, MA, USA

Jiaming Zeng
jiaming@ibm.com
IBM Research
Cambridge, MA, USA

Moninder Singh
moninder@us.ibm.com
IBM Research
Yorktown Heights, NY, USA

Marzyeh Ghassemi
mghassem@mit.edu
Massachusetts Institute of Technology
Cambridge, MA, USA
Vector Institute
Toronto, Ontario, Canada

available to ML models and human experts and the implications of model-detectable differences in clinical notes. Our work makes three key contributions. First, we find that models can identify patient self-reported race from clinical notes even when the notes are stripped of explicit indicators of race. Second, we determine

Review

Past Year Mental Disorders among Adults in the United States: Results from the 2008–2012 Mental Health Surveillance Study

Rhonda S. Karg [1], Jonaki Bose [2], Kathryn R. Batts [1], Valerie L. Forman-Hoffman [1], Dan Liao [1], Erica Hirsch [1], Michael R. Pemberton [1], Lisa J. Colpe [3], Sarra L. Hedden [2]

In: CBHSQ Data Review. Rockville (MD): Substance Abuse and Mental Health Services Administration (US); 2012.
2014 Oct.

Depression Risk


Patching in demographics can be used to study *implicit* biases.

Step 1


31 y/o ~~woman~~ presents with…

Remove all explicit mentions of gender (race) and sexed conditions

Step 2




Depression Risk



Patch residual activation of _ (chat token) in $x_{depression}$ at layer ℓ with the residual activations of the "Male" token at layer ℓ from x_{male}

Depression Risk



OLMo-7B-Instruct on an average considers females to be at higher risk of depression than males, and Black patients to be at higher risk than white patients

$$\Delta_{\text{risk}} = \frac{1}{|S|} \sum_{i=1}^{|S|} (u_i - v_i)$$

S	set of brief hospital courses
u	binary risk prediction when female (Black) is assigned
v	binary risk prediction when male (white) is assigned

Depression Risk

Male	Female
...there is no direct indication that the patient is at risk of depression....	...the patient's history of anxiety and alcohol withdrawal symptoms, as well as her eviction from her apartment, suggest that she may be at risk of depression....
...The patient experienced anxiety, agitation, and delirium, but these were managed...	...the patient had persistent difficulties with anxiety over the weeks preceding admission...

White	Black
...there is no indication that she is at risk of depression...	...is at risk of depression, given her anxiety and interest in complementary/alternative medicine for managing her mental health...
...the patient is not at risk of depression...	...at a higher risk of depression...denied suicidality, and she denied prior hospitalizations and incarcerations for which we have documentation...

Step 1

31 y/o African-American presents with…


Keep all explicit mentions of race

Step 2

Can we spot this reliance on race?

Read the below brief hospital course of a patient.

31 y/o African-American presents with…




Yes

Is the patient at risk of depression? Answer Yes or No.


We would like to reveal and mitigate undue reliance on race.


Sparse Autoencoder (SAE)



Can SAEs reveal and mitigate racial bias?

I like cats because...





- Train race probe using patient discharge summaries.
- Focus on patients who identified as “White” and “Black/African-American”.

Black predictive latent

Term “African-American” ethnicity

gemma-2b

Term “African-American” ethnicity, and medical conditions

gemma-9b

Top predictive latent description

General domain

…trends within the African American…

…in the universe of black studies…

…questions in African American history…

…now Morehouse College, where he…

…Category… Howard University..


…Black folks we'd like…

…apparently NAACP paid for it…


Top tokens

afro
African
Jamaican
african
NAACP
Afro

Clinical domain




gemma-2-2B




gemma-2-9B

Clinical domain




gemma-2-2B



gemma-2-9B

Does the Black latent have a causal effect on model output?

Given latent activates on discussions of violence, we evaluate whether steering with the Black latent induces the model to view patients as violent.



Steering with the Black latent increases the risk assigned to the probability that a patient will become "belligerent".

Prompt: *Read the following clinical note of a patient. [BHC] Based on the note, is the patient at risk of getting belligerent? Respond with 'Yes' or 'No' followed by your reasoning. You must start your answer with "Race:", followed by the patient's race.*

Before intervention	After steering
Race: Not mentioned in the note ***Answer: No*** *The note does not provide any information*	**Race: African-American** **Answer: Yes** The patient is experiencing a high level of stress and anxiety...

Model	Δ_{Black}	Δ_{white}
2B	0.51↑	-0.01
9B	0.81↑	0.09

None of the reasoning chains reveal reliance on race

$\Delta_{Black/white}$ indicates *change in positive rate for patient belligerence after steering with race latents.*

Your project!

Behavior

Interpretability method

Dataset

Behavior	Interpretability method	Dataset
Demographic bias	Activation patching, SAE	MIMIC BHCs

MedRedFlag: Investigating how LLMs Redirect Misconceptions in Real-World Health Communication

Sraavya Sambara *
Independent Researcher
srsambara@gmail.com

Yuan Pu *
Duke University
yuan.pu@duke.edu

Ayman Ali
Duke University
ayman.ali@duke.edu

Vishala Mishra
Duke University
vishala.mishra@duke.edu

Lionel Wong
Stanford University
liowong@stanford.edu

Monica Agrawal
Duke University
monica.agrawal@duke.edu

Abstract

Real-world health questions from patients often unintentionally embed false assumptions or premises. In such cases, safe medical communication typically involves *redirection*: addressing the implicit misconception and then responding to the underlying patient context, rather than the original question. While large language models (LLMs) are increasingly being used by lay users for medical advice, they have not yet been tested for this crucial competency. Therefore, in this work, we investigate how LLMs react to false premises embedded within real-world health questions. We develop a semi-automated pipeline to curate MedRedFlag, a dataset of 1100+ questions sourced from Reddit that require redirection. We then systematically compare responses from state-of-the-art LLMs to those from clinicians. Our analysis reveals that LLMs often fail to redirect problematic questions, even when the problematic premise is detected, and provide answers that could lead to suboptimal medical decision making. Our benchmark and results reveal a novel and substantial gap in how LLMs perform under the conditions of real-world health communication, highlighting critical safety concerns for patient-facing medical AI systems. Code and data are available at https://github.com/srsambara-1/MedRedFlag.




Figure 1: MedRedFlag contains patient questions with *false underlying assumptions* that human clinicians choose to *redirect* when answering. LLMs often *accommodate* false assumptions when answering instead.




Figure 1: `MedRedFlag` contains patient questions with *false underlying assumptions* that human clinicians choose to *redirect* when answering. LLMs often *accommodate* false assumptions when answering instead.

Behavior

Dataset




Figure 3: **Anatomy of a representative LLM response** to patient question with embedded false assumptions. We find that even when LLMs **address false or unsafe assumptions** in the patient question (*green*), they still often extensively **accommodate the false assumption** *(red)* with detailed, unsafe advice based on the patient question.

Interpretability method

LLMs Encode Harmfulness and Refusal Separately

Jiachen Zhao
Northeastern University

Jing Huang
Stanford University

Zhengxuan Wu
Stanford University


David Bau
Northeastern University

Weiyan Shi
Northeastern University

Abstract

LLMs are trained to refuse harmful instructions, but do they truly understand harmfulness beyond just refusing? Prior work has shown that LLMs' refusal behaviors can be mediated by a one-dimensional subspace, i.e., a refusal direction. In this work, we identify a new dimension to analyze safety mechanisms in LLMs, i.e., harmfulness, which is encoded internally as a separate concept from refusal. And there exists a harmfulness direction that is distinct from the refusal direction. As causal evidence, steering along the harmfulness direction can lead LLMs to interpret harmless instructions as harmful, but steering along the refusal direction tends to elicit refusal responses directly without reversing the model's judgment on harmfulness. Furthermore, using our identified harmfulness concept, we find that certain jailbreak methods work by reducing the refusal signals without suppressing the model's internal belief of harmfulness. We also find that adversarially fine-tuning models to accept harmful instructions has minimal impact on the model's internal belief of harmfulness. These insights lead to a practical safety application: The model's latent harmfulness representation can serve as an intrinsic safeguard (*Latent Guard*) for detecting unsafe inputs and reducing over-refusals that is robust to finetuning attacks. For instance, our *Latent Guard* achieves performance comparable to or better than *Llama Guard 3 8B*, a dedicated finetuned safeguard model, across different jailbreak methods. Our findings suggest that LLMs' internal understanding of harmfulness is more robust than their refusal decision to diverse input instructions, offering a new perspective to study AI safety[1].

Data



MIMIC-IV-Core

Patient demographics, admission tracking, and stay information is available in MIMIC-Core.

Read more ...

MIMIC-IV-Hosp

The Hosp module provides all data acquired from the hospital wide electronic health record. This includes laboratory measurements, microbiology cultures, medication information, services provided, billed diagnoses and procedures, and so on.

Read more ...

MIMIC-IV-ICU

The ICU module contains information collected from the clinical information system used within the ICU. This includes highly granular information such as hour-to-hour vital signs, information about fluid management, and other charted observations.

Read more ...

MIMIC-IV-ED

The ED module contains data for emergency department patients including a triage assessment, nurse-validated vital signs, medicine reconciliation, and treatment information.

Read more ...

MIMIC-IV-CXR

MIMIC-cxr provides chest x-ray images and radiology reports for a subset of patients admitted to the emergency department.

Read more ...

MIMIC-IV-Note

The Note module contains deidentified free-text clinical notes for hospitalized patients.

Read more ...

MIMIC-III

MIMIC-III is an older version of MIMIC. It contains an older group of patients (ending in 2012), and a subset of the ICU and hospital information available in MIMIC-IV. We highly recommend researchers starting new studies to use the above modules in MIMIC-IV. We will maintain MIMIC-III for a short period to allow researchers to transition.

Read more ...